

Production models

fish610.055 Fisheries and ecosystem models

Gunnar Stefansson

April 10, 2018

History

Stock-production models: Typical for whales, shrimp, nephrops and other difficult to age species

Originally: Assumed equilibrium (oversimplified)

Now: Use dynamic stock-production models

An equilibrium model

A simple equilibrium production model:

$$Y = rB\left(1 - \frac{B}{K}\right)$$




Figure : Equilibrium yield and biomass

Simple forward projections

Given B the production is $rB(1 - B/K)$.

If the catch is Y the stock biomass will become

$$B + rB(1 - B/K) - Y.$$

Notably, the stock increases if Y is less than the production and vice versa.




Figure : Equilibrium yield with a constant catch line.

The problem with equilibrium models

Over-fished=catch above sustainable yield

Observed biomass and catch all above the underlying curve

Note: Will overestimate production if assume equilibrium




Figure : The bias issue when estimating data from an equilibrium model. The solid line is the true model and the dot/dashed line is from generated data when fishing is 1.5 times the equilibrium catch.

A dynamic biomass model

A simple forward projection model:

$$B_{y+1} = B_y + rB_y(1 - B_y/K) - Y_y$$

where the catch is specified according to some rule, e.g. proportional harvest:

$$Y_y = pB_y.$$




Figure : Equilibrium yield and curve describing the trajectory of catch and biomass under a constant harvest strategy.

Unknown parameters

Model: $B_{y+1} = B_y + rB_y(1 - B_y/K) - Y_y$

Unknown parameters: r , K , and B_0 .

- Given model and parameters, the trajectory can be generated.
- Assumes catches are known constants (no error)
- Parsimony: May (need to) assume $B_0 = K$.