

Data vectors

math612.1 612.1 Numbers, arithmetic and algebra

Gunnar Stefansson (editor) with contributions from very many students

September 3, 2018

The plane

Pairs of numbers can be depicted as points on a plane.
The plane is normally denoted by \mathbb{R}^2 .

Simple plots in R

Graphing functions in R

- `plot` - plots a scatter plot (as a line plot)
- `points` - adds points to a plot
- `text` - adds text to a plot
- `lines` - adds lines to a plot




Figure : Points on a plane, drawn with R.

The following R commands can be used to generate a plot with two points:

```
> plot(1,2,xlim=c(0,5),ylim=c(0,5),xlab="x",ylab="y")
> points(3,1)
> text(1,2,"(1,2)",pos=4, cex=2)
> text(3,1,"(3,1)",pos=4, cex=2)
```

Data

Data are usually a sequence of numbers, typically called a vector.

Indices for a data vector

If data are in a vector x , then we use indices to refer to individual elements. If i is an integer then x_i denotes the i 'th element of x . Note that we do not distinguish (much) between rows and columns.

Summation

We use the symbol Σ to denote sums.

In R, the sum function adds numbers.

If $x = (4, 5, 3, 7)$ then

$$\sum_{i=2}^4 x_i = x_2 + x_3 + x_4 = 5 + 3 + 7 = 15.$$

Within R:

```
> x<-c(4,5,3,7)
> x
[1] 4 5 3 7
> sum(x)
[1] 19
>
```