

# Inverse functions and the logarithm

math612.1 612.1 Numbers, arithmetic and algebra

Gunnar Stefansson (editor) with contributions from very many students

September 3, 2018

# Inverse Function


If  $f$  is a function, then the function  $g$  is the inverse function of  $f$  if

$$g(f(x)) = x$$

for all  $x$  in which  $f(x)$  can be calculated

# When the inverse exists: The domain question

Inverses do not always exist. For an inverse of  $f$  to exist,  $f$  must be one-to-one, i.e. for each  $x$ ,  $f(x)$  must be unique.



**Figure :** The function  $f(x) = x^2$  does not have an inverse since  $f(x)=1$  has two possible solutions -1 and 1.

# The base 10 logarithm

When  $x$  is a positive real number in  $x = 10^y$ ,  $y$  is referred to as the base 10 logarithm of  $x$  and is written as:

$$y = \log_{10}(x)$$

or

$$y = \log(x)$$

# The natural logarithm


A logarithm with  $e$  as a base is referred to as the natural logarithm and is denoted as  $\ln$  :

$$y = \ln(x)$$

if

$$x = e^y = \exp(y)$$

Note that  $\ln$  is the inverse of  $\exp$ .



**Figure :** The curve depicts the function  $y = \ln(x)$  and shows that  $\ln$  is the inverse of  $\exp$ . Note that  $\ln(1) = 0$  and when  $y = 0$  then  $e^0 = 1$ .

# Properties of logarithm(s)

Logarithms transform multiplicative models into additive models, i.e.

$$\ln(a \cdot b) = \ln a + \ln b$$

# The exponential function and the logarithm

The exponential function and the logarithms are inverses of each other

$$x = e^y \Leftrightarrow y = \ln x$$